

Research

Open Access

Geographical and seasonal correlation of multiple sclerosis to sporadic schizophrenia

Markus Fritzsche*

Address: Clinic for Internal Medicine, Soodstrasse 13, 8134 Adliswil, Switzerland

Email: Markus Fritzsche* - markus.fritzsche@bluewin.ch

* Corresponding author

Published: 20 December 2002

Received: 28 November 2002

International Journal of Health Geographics 2002, 1:5

Accepted: 20 December 2002

This article is available from: <http://www.ij-healthgeographics.com/content/1/1/5>

© 2002 Fritzsche; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

Abstract

Background: Clusters by season and locality reveal a striking epidemiological overlap between sporadic schizophrenia and multiple sclerosis (MS). As the birth excesses of those individuals who later in life develop schizophrenia mirror the seasonal distribution of Ixodid ticks, a meta analysis has been performed between all neuropsychiatric birth excesses including MS and the epidemiology of spirochaetal infectious diseases.

Results: The prevalence of MS and schizophrenic birth excesses entirely spares the tropical belt where human treponematoses are endemic, whereas in more temperate climates infection rates of *Borrelia garinii* in ticks collected from seabirds match the global geographic distribution of MS. If the seasonal fluctuations of Lyme borreliosis in Europe are taken into account, the birth excesses of MS and those of schizophrenia are nine months apart, reflecting the activity of *Ixodes ricinus* at the time of embryonic implantation and birth. In America, this nine months' shift between MS and schizophrenic births is also reflected by the periodicity of *Borrelia burgdorferi* transmitting *Ixodes pacificus* ticks along the West Coast and the periodicity of *Ixodes scapularis* along the East Coast. With respect to Ixodid tick activity, amongst the neuropsychiatric birth excesses only amyotrophic lateral sclerosis (ALS) shows a similar seasonal trend.

Conclusion: It cannot be excluded at present that maternal infection by *Borrelia burgdorferi* poses a risk to the unborn. The seasonal and geographical overlap between schizophrenia, MS and neuroborreliosis rather emphasises a causal relation that derives from exposure to a flagellar virulence factor at conception and delivery. It is hoped that the pathogenic correlation of spirochaetal virulence to temperature and heat shock proteins (HSP) might encourage a new direction of research in molecular epidemiology.

Background

Schizophrenia and multiple sclerosis are distinct neuropsychiatric disorders of the central nervous system (CNS). Schizophrenia is characterised by disturbances in multiple domains of brain functioning, few *in vivo* or *post-mortem* studies of which find evidence for a particular structural alteration [1]. MS typically manifests as acute focal inflammatory demyelination and axonal loss in-

volving the immune system and culminating in the chronic multifocal sclerotic plaques from which the disease gets its name [2-4]. MS and schizophrenia, nonetheless, exhibit a striking epidemiological overlap [5]. Prevailing in the colder parts of the world and affecting mainly young adults in their most productive years, both run an irregular, chronic course. Apart from acute infections [6], no other disease exhibits an equally marked ep-

idemiological cluster by season and locality, nurturing the hope that solutions might ultimately be attainable [5].

Geographical correlations between MS and schizophrenia prevalence rates (PRs) have been reported worldwide [5,7–9]. In the north of the USA, the states with the highest rates of schizophrenia score significantly higher rates of MS than the states with the lowest schizophrenia rates in the south [9]. Findings that immigrants from tropical low-risk areas are more likely to contract schizophrenia or MS than those who stay behind have been widely replicated and are now considered strong epidemiological risk factors (for review see [10–12]). Migration away from the hot spots reduces the probability of developing MS if the move takes place before adolescence. Regardless of the direction of the move, however, MS death rates for migrants born in one risk area and dying in another are intermediate between those of their birthplace and their final residence. These biological gradients suggest a common environmental component that could be influenced. If caused by an infectious agent [12] as presupposed by Marie in 1884 [13], what kind of transmission or virulence would lie at its root?

The earliest lesion seen in MS is a focal infiltration of lymphocytes around small blood vessels in the brain and spinal cord. This implies that inflammatory cells are reacting against an antigen located in the central nervous system (CNS) to which they have become sensitised. In MS patients, the intrathecal synthesis of immunoglobulins, the appearance of immune complexes and change in the balance of T cell populations all indicate immune activity in tissues that are normally quiet (for review see [2,3]). The responsible antigen could be CNS tissue ('self'), in which case we are dealing with an autoimmune disease. Alternatively, it could be of microbial ('foreign') origin, or both. Analogous to streptococcal infection in relation to rheumatic fever, the environmental factor in MS may be accommodated by arguing that exposure to an infection elicits upon re-exposure an abnormal immune response against a similar antigen in the CNS.

From this epidemiological point of view, we are best advised to search for positive and negative correlation. For general immunity acquired against endemic infection is supposed to increase the resistance against MS in developing countries [14]. In the interior of New Guinea, in fact, where neither MS [15] nor schizophrenia [16] nor Lyme borreliosis [17] appears to be prevalent, the inhabitants exhibit antibodies possibly induced by endemic treponemes that cross-react with *Borrelia burgdorferi* antigens [17].

Since the beginning of the last century it was suggested that MS and neurosyphilis caused by the spirochaete

Treponema pallidum [18] had similar clinical and histological characteristics. Extremely difficult to find, as in tabes dorsalis [18], neuropathologists documented the presence of spirochaetal structures in MS plaques suggesting that the patients were infected with a spirochaete [19,20], most likely *B. burgdorferi* [21].

Congenital infection by *B. burgdorferi* resembles congenital syphilis as well. Like in humans [22], however, chronic prenatal infections by *B. burgdorferi* are rare in mice, and acute prenatal infection of the unborn is restricted to a narrow time window of transplacental transmission [23]. Henceforth the birth excess of those individuals who later in life develop sporadic schizophrenia mirrors the seasonal distribution of *Ixodes* ticks and *Borrelia burgdorferi* at the time of conception [16].

If in analogy to chronic hepatitis B infection at birth [24], MS were induced [25] or exacerbated [26] by exposure to *B. burgdorferi* antigens during delivery, we would expect a direct seasonal match between MS birth excesses and Ixodid tick activity worldwide. Contrasting with the possible deleterious mutagenic effect of *B. burgdorferi* at conception [16], a time difference of nine months is expected between MS births and the schizophrenic insult to the implanting embryo. Should general immunity acquired against treponemes [17] increase resistance against *Borrelia* and MS, we would furthermore expect a positive geographic correlation of MS to neuroborreliosis as well as a negative correlation of MS to endemic treponematoses. Otherwise, the present hypothesis of *B. burgdorferi* as a possible major aetiologic factor for both congenital sporadic schizophrenia and MS would be falsified.

Methods

The different prevalence rates of MS as recently reviewed [15] were compared to the global distribution of endemic treponematoses [27] and to all significant schizophrenic birth-excess rates [16] worldwide. For statistical reasons, only publications encompassing more than 3000 cases of significant schizophrenic birth excesses compared to the normal population have been considered [28], including two studies from Denmark and Australia showing both one significant and one non-significant result each. In a second step of the investigation, the routes of *Borrelia* carrying migratory birds were considered for the following reason. The schizophrenic birth excesses are limited to exactly those regions that are endemic for *B. burgdorferi* transmitting *Ixodes* tick vectors [16]. This geographical overlap, however, is less than perfect with respect to MS, particularly in southern latitudes, and the use of artificial country boundaries is of doubtful biological value, unless they are given by naturally defined geographical areas [11]. As the rule of MS varying with latitude is violated in central Europe including Switzerland and its neighbour-

ing countries, areas of high relative risk were put under closer scrutiny. MS hot spots in central Europe, in fact, harbour the few remaining nesting sites of the classical passerine bird, the white stork (*Ciconia ciconia*), and migratory seabirds following the rivers and islands spread *Ixodes* ticks and *Borrelia garinii* (a subspecies of *Borrelia burgdorferi sensu lato*) worldwide [29] (see Figure 1). From the comprehensive literature, the seasonal periodicity of Ixodid tick activity was then used in this meta analysis and plotted against numerical data encompassing all neuropsychiatric birth excesses [30] of the respective macroclimatic regions including MS (Figure 2).

Results

Geographical correlation of MS and sporadic schizophrenia to *Ixodes* ticks and migratory birds

The geographical gradient of MS (Figure 1), which sharply declines at the 37° latitude [12,15], entirely spares the tropical belt where human treponematoses caused by *T. carateum*, *T. pertenue*, and endemic *T. pallidum* prevail [27]. This negative correlation between areas of endemic treponematoses on the one hand and MS as well as schizophrenic birth excesses on the other is striking. With the notable exception of Florida, Australia and Tunisia (see Figure 1), subtropical areas exhibit neither a significant schizophrenic birth excess nor a prevalence rate (PR) of MS higher than 5/100'000 [15].

In central Europe, the highest relative risks for MS can be found along the breeding sites of migratory birds. These hot spots are located along the north-south axis of the Upper Rhine plain and its tributaries, the Bas Rhin, Haut Rhin, Moselle regions [31], around Basel [32], in the Swabian Alps [33], Sardinia, as well as the Aosta Valley [15], the only place in Italy where the white stork (*Ciconia ciconia*) still nests (see insert in Figure 1). Tunisia, which is reached by passerine European birds carrying *Ixodes* ticks and *B. garinii* [34], scores the highest rate of MS in Africa [15]. Malta, by contrast (see Figure 1), which compared to Sicily is relatively free of MS [11,15], hosts a number of endemic lizards, and ticks lose their infectious potential for humans when feeding on these reptiles. This so-called zooprophyllactic effect also applies to the low-risk areas of MS and sporadic schizophrenia [9] in the United States, where south of the 37° latitude infections by *B. burgdorferi* are poorly maintained in lizards [35].

Although of low prevalence, MS exists in South East Asia: in Japan and Taiwan [15,36] down to the Philippines [12], where the Wallace Line (see Figure 1) limits the southward spread of *B. burgdorferi* harbouring Ixodid ticks into Austronesia [37]. In Irian Jaya (I.J.) and P.N.G. (see Figure 1), from where neither the presence of Lyme disease nor MS has been reported, schizophrenia is non-existent apart from the south-west coast which is

sporadically reached by migratory birds and ticks [16]. In the southern hemisphere, MS [15] and schizophrenic birth excesses [38] are, compared to the northern hemisphere, less significant. From the area of Singapore, which is non-endemic for *Ixodes* ticks and *B. burgdorferi* but endemic for treponematoses (see Figure 1), a schizophrenic birth excess is significantly absent. This trend parallels the scarcity of MS and relatively recent upsurge of schizophrenic birth excesses in Japan, where *B. garinii* harbouring ticks have been sporadically introduced by migratory birds from North East Asia (for discussion see [16]). Southern Australia and New Zealand, by contrast, which can be reached by polar seabirds carrying *Ixodes uriae* and *B. garinii* via the Antarctic [39], score relatively high rates of schizophrenia [40] and MS [15].

Geographical correlation of MS to *Borrelia burgdorferi sensu lato*

The infection rates of *B. garinii* in *I. uriae* ticks [29] reflect not only the global distribution of MS, but also its worldwide gradient. The highest number of spirochaetes detected by microscopy were collected from seabirds in Iceland, Alaska and the Faroes (see Figure 1) all areas from where epidemics as well as very high rates of MS have been published [12,41]. Ticks from Sweden yielded lower spirochaetal counts and on an island south of New Zealand, where MS is reportedly less prevalent compared to the northern hemisphere [15], the number of detected spirochaetes was the lowest [29]. Ticks from Cape Sizon (France) yielded negative results, and, unfortunately, the ticks collected on the Falklands and the Crozet Islands (South Africa) were dead and thus not examined by microscopy. Regardless of the geographical origin, *B. garinii* DNA was isolated and detected by PCR in all ticks and cultured spirochaetes, with two notable exceptions. The samples from the Atlantic coast of France and those from the Falklands - which do not appear as particularly hot spots for MS [31,42] - were all negative [29].

Seasonal correlation of MS and sporadic schizophrenia to *Ixodes* tick activity

If the stochastic annual fluctuations of schizophrenic births [43] and Lyme disease [44] are taken into account, the birth excesses of MS in Denmark [8] and those of schizophrenia in Finland [45] are exactly nine months apart. In northern Europe, the seasonal patterns thereby reflect the activity of endemic *Ixodes ricinus* [44,46] at the time of conception and parturition respectively. Likewise, there exists a nine months' shift between MS birth excesses in Canada near Vancouver [30,47] and schizophrenic births in the USA [48]. The respective seasonal distribution exactly mirrors the periodicity of the adult and juvenile stages of *B. burgdorferi* transmitting *I. pacificus* ticks along the West Coast [49] and that of *I. scapularis* ticks along the East Coast (see [37]). Amongst all other neu-




Figure 1
Geographical correlation of schizophrenic birth excesses and MS prevalence to spirochaetal diseases The gradient of MS prevalence [15] and schizophrenic birth excesses [16,28] entirely spares the tropical belt where human treponematoses prevail [27]. In subtropical zones between the 37° latitudes, there exists an additional climatic effect diminishing the prevalence rates of MS worldwide. This 'rule' of variation by latitude, however, is violated in Europe, Australia and New Zealand where circumpolar migratory seabirds reportedly introduce *Ixodes uriae* and *Borrelia garinii* [29] from the MS hot spots in the north [12,15]. The arrows represent the migratory routes and distribution of seabirds if a number of species such as puffins (shearwaters), seagulls and terns are taken together [56–59]. The sites from where samples in search for *Borrelia garinii* were collected by Olsen et al. [29] are marked with a red dot and include the Egg and St. Lazaria Islands (Alaska), Gannet Island in Newfoundland (Canada), Iceland, the Falklands, the Faroes, Bonden Island (Sweden), Cape Sizun (France), the Crozet Islands (South Africa), and Campbell Island (New Zealand). There are just a few subtropical areas including Madeira, Morocco and Tunisia from where the presence of *B. burgdorferi s.l.* has been documented. Two protective factors could thus explain the rarity of MS and schizophrenic birth excesses in developing countries: acquired immunologic resistance against spirochaetes in the tropics, and higher temperature being inversely related to tick-borne spirochaetal transmission in subtropical zones.

ropsychiatric disorders, only amyotrophic lateral sclerosis (ALS) [50] shows a similar seasonal trend [30,51] with respect to *Ixodes* tick activity in Europe [37] (see Figure 2).

Discussion

The uneven distribution of MS has been noted in the medical literature for well over 100 years [52]. Although difficult to assess reliably from one geographical area to another [11], it has also become clear that the frequency of the disease varies significantly not only in different parts of the world, but even within countries. The first epidemiological study by Davenport (1922) pointed out that MS affected persons of Scandinavian and Finnish descent more than other ethnic groups, a conclusion confirmed by Bailey's study of American troops in World War

I. Later propositions that MS, as well as ALS [53,54], relate to latitude and cold climate unleashed a controversy over nature versus nurture that continues to this day (for review see [11,12]).

Migratory seabirds spread *B. burgdorferi sensu lato* (*B. garinii*) worldwide

Since current opinion favours a genetic/immune process for MS, as well as ALS, the possibility of spirochaetal infection has not been considered in depth. The similar global distribution of seabird-borne ticks and Scandinavians and their descendants, who appear to be at high genetic risk for MS [55], is not entirely coincidental. One Viking legend claims that the Faroes were discovered by following marine birds [56]. While seabirds themselves often follow




Figure 2

Seasonal correlation of neuropsychiatric birth excesses to *Ixodes* ticks and Lyme borreliosis in America and Europe In America, the seasonal distribution of schizophrenic [48] and MS [47] birth excesses exactly mirrors the periodicity of *Ixodes scapularis* along the East Coast [37] and that of *Ixodes pacificus* along the West Coast [49]. Curiously enough, in many studies conducted in the USA, the month of June is often the month with the lowest risk of developing schizophrenia (Jean-Paul Selten, personal communication, 2002). This schizophrenic birth deficit corresponds to the tick activity, which is at its lowest nine months earlier in September, being particularly striking, if both species *Ixodes scapularis* and *Ixodes pacificus* are taken together. If stochastic annual fluctuations are taken into consideration [43,44], the spring (a) population of *Ixodes ricinus* [37] likewise mirrors the significant birth excess rates of schizophrenia [45], ALS [51] and MS [8] in Europe. The nine months' shift between sporadic schizophrenia on the one hand and ALS and MS on the other reflects the possible transplacental transmission of *Borrelia burgdorferi* [23] at the time of conception [16] and delivery [24].

ships and eat fish offal floating on the water surface [57], the Scandinavians were apparently led by such birds to new fishing grounds [56].

Little more than 300 of the approximately 9000 known bird species are seabirds. Yet, these birds occur throughout the world and are the only group of birds to have successfully colonised Antarctica, the most inhospitable continent. Although some seabirds breed and winter within comparatively small areas, many are champions in long distance migration, travelling thousands of kilometres on journeys from the far north of the northern hemisphere to the limits of the Antarctic pack ice in the southern hemisphere [57,58].

The seabirds, which spend most of their lives far out to sea, must procreate on land. To avoid predators, they often breed on isolated islands and peninsulas leading to the aggregation of hundreds of thousands, and sometimes millions of pairs during the breeding season. The presence of such huge numbers not only depends on the abundance of food in the surrounding seas, the crowding habit

also makes the birds and their offspring vulnerable to the seabird associated *Ixodes uriae* tick. No wonder, infestation by *I. uriae* has been reported from more than 50 species of seabirds in both hemispheres (see [29]).

Because seabirds take their food from shallow waters, many coastlines and islands are prime sites for breeding colonies [57,58], *Borrelia* transmitting *I. uriae* [29], and MS [56]. On the Faroe Islands (see Figure 1), where *Ixodes uriae* reportedly transmits *Borrelia* from seabirds to human bird catchers [59], the MS scenario apparently unfolded after an annulled ban on fowling seabirds during a food shortage in World War II [56]. Whatever speculation be more plausible - transmission by ticks [56] or soldiers [12,14] - the prevalence of MS subsequently rose from apparently zero to 21 cases heralding the first of four successive epidemics [12].

On Iceland and elsewhere such epidemics have been attributed to increased awareness, changes in ascertainment or better diagnosis of MS, particularly of more benign cases in the post-war era [11]. However, a common setting

for MS 'epidemics' is proximity to coastal areas or islands where seabirds nest [56]. From western Alaska both the presence of *Ixodes uriae* and *B. garinii* have been reported [29], and in Sitka being surrounded by three major seabird colonies in south-eastern Alaska (see Figure 1) MS was unknown until its first outbreak occurred in 1965 [41].

Genetic [60,61] as well as epidemiological studies, in which biological plausibility had been ignored [11], often provided contradictory information. One of the significant risk factors mentioned in a study from Key West (Florida), for instance [62], were visits to a local military base, a finding that the authors noted as a point of similarity to the military occupation and its reported effect on the Faroe Islanders [12]. In Malta, however, which was occupied by British troops from 1802 to 1978, the low PR of MS doubled from 1978 to 1988 (from 4.2 to 8.4), after the British left [11]. In contrast to Sicily with a relatively high MS PR of 61, however, the population of Malta (see Figure 1) still enjoys significant protection from MS that cannot be convincingly explained by geneticists either. It is noteworthy that since the Arabs were driven out by a band of Scandinavian adventurers, who had established a kingdom in southern Italy, Malta became a Norman appendage of Sicily for almost half a millennium. Yet, in addition to Scandinavian genes [55], Malta also hosts four endemic races of lizards (*Podarcis filfolensis*), and ticks lose their infectious potential for human beings when feeding on these reptiles. This so-called zoophylactic effect also applies to the USA [35], where south of the 37° latitude (see Figure 1) the PR of MS is significantly lower [12,15]. The only notable exception is Florida [62], where migratory seabirds stop over for nesting [56]. Otherwise, infections by *B. burgdorferi* are poorly maintained by lizards in the south of the United States [35]. While seabirds are not common in states such as North Dakota, Montana, Idaho and Colorado where MS is high, the inland spread of *Ixodes* ticks and Lyme disease by land birds is well documented in the United States [63].

At the expense of a persistent vulnerability to lizards, the tick-borne pathogen has proven successful in spreading neuroborreliosis including MS via sub-polar routes [29] across the globe (see Figure 1). European strains of *B. garinii* infections have been documented in human cases from the USA [64] and Australia [39]. The case of Lyme borreliosis in the southern hemisphere has been confirmed by culture and serotyping [39]. Being of European origin, this type of *B. burgdorferi* s.l. was most probably introduced into Australia by a migratory seabird. Not surprisingly after all, southern Australia and New Zealand, which polar seabirds carrying *Ixodes uriae* [29] reach via the Antarctic, score relatively high risks of MS. Even the highest PR in these communities, largely originating from

the United Kingdom, is not much more than half the rate in most parts of the British Isles [15]. This difference in relative risk is hard to understand from a purely genetic point of view. But there isn't much room for 'pure' environmentalists either, as Waikato in New Zealand, where the main step in MS morbidity occurs across the North Island, scores a lower rate than places in Australia on a comparable southerly latitude [2,15]. For migratory seabirds introducing *Ixodes uriae* from the northern hemisphere [29] reach New Zealand later than Australia (see Figure 1).

In the northern and southern hemispheres, several species are responsible for this transhemispheric exchange. Great puffins or Manx shearwaters (*Puffini puffini*), for example, move around the world in giant loops. They are abundant off the European continental shelf in July and August, when they are heading southeast. Between September and December, the puffins spend their time mainly along the American coast from Rio de Janeiro in the north to the Rio de la Plata in the south. Along these coasts, the nutrient-rich water advances during this period with the Falkland Current producing upwellings on the water surface rich in fish. By March and April the birds leave their breeding colonies on the Falklands and other islands in the South Atlantic heading northwest across the equator to the rich fishing waters off Newfoundland. Then they gradually move back across the North Atlantic, where they are often seen around Scotland, Ireland and the Faroes, where the traditional puffin-hunting season starts in the end of July [57–59].

In the southern oceans, where the winds blow almost continuously eastwards in the roaring forties and furious fifties, a ringed great puffin has even been found in south Australia. The distribution of short-tailed puffins, or short tailed shearwaters (*Puffini tenuirostri*), is limited to this part of southern hemisphere, where the birds breed on islands off the coast of New Zealand and Australia. In Tasmania, as in the Faroes, their so-called mutton-bird chicks are regularly fowled [58].

Among the most successful and widespread marine birds are the seagulls. There are some 45 species, which occur in both hemispheres. Colonies may be tens of thousands strong, particularly if there is a major source of food thereby. Outside the breeding season most gulls, such as the black headed gull (*Larus ridibundus*), undertake migratory movements, sometimes wintering well out to sea. They move parallel to latitude to avoid cold weather, or they simply disperse over comparatively short distances along rivers [58,65] (see Figure 1). Along the tributaries of the upper Rhine [65] they thus reach the foot of the Swiss Alps where *B. garinii* has become highly endemic among other terrestrial passerine birds. These, in turn, may spread borreliosis to other hosts [66] by regularly migrating to north-

ern [67] and southern Europe either along the valley of the upper Rhone or directly via Alpine passes (personal communication 2002, Christian Marti, Swiss Ornithological Institute, Sempach).

The seasonal correlation of MS to Ixodes ticks may explain hitherto discrepant findings

Extending a previous report from the United States, Timpler and colleagues found a high geographical correlation between MS and schizophrenia in Italy [7]. As a correct temporal relation between cause and effect is essential in epidemiology [6], correlated birth patterns of MS and schizophrenia were then studied. In Denmark, a significant birth excess of MS was disclosed in spring-early summer, but the data on schizophrenia were insignificant compared to the general population [8]. This negative finding, which can be explained by stochastic fluctuations (for discussion see [43]), was unfortunate since accumulating evidence from most other studies [28] did yield significant schizophrenic birth excesses in winter and spring.

In Sicily, interestingly enough, MS birth excesses being shifted towards the end of the year [68] show a seasonal trend reminiscent of the seasonal *Ixodes* tick activity in neighbouring North Africa [37]. And in Tunisia, where *Ixodes* ticks harbour *B. garinii* and *B. lusitaniae* [34] - species known to be scattered geographically by birds from Europe - MS scores the highest rates in Africa [15].

Temperature and spirochaetal virulence: the genetic interface between immunity and environment

MS has never been reported in ethnically pure Eskimos, Inuit, Lapps, Amerindians, Australian aborigines, New Zealand Maoris or Pacific Islanders [11]. Yet, most of these natives either live near the polar circle, where tick activity abates due to low ambient temperature, or they live in tropical climates where *Treponema* [27] but not *Borrelia* spirochaetes prevail (see Figure 1).

The most convincing evidence for the importance of genetic or acquired resistance is the extreme rarity of MS in native Africans. In the Cape Province of South Africa, the disease is recognised among the so-called coloured, but neurologists in Johannesburg are extremely reluctant to make the diagnosis in a black person [11]. Hawkes [14] maintains that in developing countries a general immunity is acquired against infection, which might also spread "in utero or during parturition" and possibly increases the resistance against MS and sexually transmitted disease. Until the early nineties, I was working as a medical delegate in southern Africa and cannot share Hawkes's opinion on the isolation of 'black' Africans from 'white' sexual permissiveness. Venereal diseases including syphilis were highly prevalent among natives of South Africa, Angola and Mozambique, and in that part of the world it is the in-

cidence of AIDS, but not MS, which has substantially risen. The "infrequency of MS and AIDS in the same patient" [14] rather favours the hypothesis of a hyperergic immune process [2,3] which is suppressed in a state of immunodeficiency such as AIDS.

Whilst it is always more prudent to await the discovery of new facts [69], long-held [25] hypotheses [14] often raise testable questions if taken together and dissected with Occam's razor. The MS gradient, which sharply declines at the 37° latitude, suggests a temperature-related environmental factor that cannot be ignored (see Figure 1). The geographical distribution of endemic treponematoses, by contrast, as well as blood group O antigens which appear to convey resistance against *Treponema pallidum* [70], are restricted to exactly those parts of the tropics that are free of MS. This contrast is of evolutionary importance with regard to the adaptation of intracellular pathogenic spirochaetes. Like the ABO blood group system [71], the evolutionary conserved heat shock proteins (HSPs) do not only induce heat resistance, but also activate host immune defences that are detrimental for pathogens (see for example [72-78]).

Molecular evidence reveals that *Treponema pallidum* - the agent of syphilis being an exclusively human pathogen and *B. burgdorferi* s. l. - the human and animal pathogens of Lyme borreliosis - have circumvented this immunological impasse differently. Whilst in the course of evolution *Treponema pallidum*, being directly transmissible from human to human, lost its capacity to induce HSPs [79], heat resistance [80] and thus vector-borne transmission, *B. garinii* has adapted to a broad temperature range [81]. To survive heat shocks during vector-borne transmission, the gram-negative *Borrelia* pathogen therefore expresses HSP-60 and HSP-70 [82], which are members of the evolutionary conserved HSP family. This form of vector adaptation is essential during rapid changes in temperature, in particular when transmitted from ticks to their warm-blooded hosts [83] including birds.

In the absence of other vertebrate hosts on certain islands, the presence of *Borrelia* in *I. uriae* ticks suggests seabirds to be competent reservoirs and amplifying hosts. In contrast to *B. burgdorferi sensu stricto* being cleared from the respective *Ixodes* vectors at 37°C [84], a temperature of 38°C is permissive for the transmission of *B. garinii* [81]. The relatively low body temperature of 38°C in marine birds, compared to the body temperature of terrestrial birds of 40°C, explains why these *B. burgdorferi* s.l. spirochaetes are particularly adapted to seabirds [29].

But the pathogen's success of transmission also depends on its ability to replicate and survive within a host for long periods. One option is to remain latent inside the long-

Table 1: Possible molecular mimicry between *Borrelia burgdorferi* fbrp and human interleukin-1 receptor antagonist

```

fbrp:    106 GYVELPNVNLVEEMVDMI 123
          GY++ PNVNL EE +D++
Il-1ra: 34 GYLQGNVNL-EEKIDVV 50

```

The occurrence of homologies between *B. burgdorferi* flagellae and host molecules has been amply demonstrated [95,96]. This phenomenon, termed molecular mimicry, explains why an antibody generated against a particular epitope of an infectious pathogen may become an autoantibody, reacting with a homologous epitope in the host and bringing about structural dysfunction or tissue damage. The autoimmune response will continue to attack the host even after clearance of the pathogen from the infected organism. However, autoimmunity by molecular mimicry will only occur if epitopes of the pathogen and of the host are similar enough to allow immunological cross reaction, yet different enough to break immunological tolerance [101]. The homology was assessed by protein BLAST using the search program at OMIM [99] for short, nearly exact matches. Length = 153, Score = 27.8 bits (58), Expect = 15, Identities = 10/18 (55%), Positives = 15/18 (82%), Gaps = 1/18 (5%). Accession numbers: *B. burgdorferi* fbrp: I448943; Human interleukin-1 receptor antagonist: 999512.

lived cells of the CNS whose temperature is about 38° in humans. This coincidence elucidates the characteristic spread and neurotropism of *B. garinii*, which is frequently associated with neurological manifestations [85]. *In vitro* evidence suggests early invasion of the CNS by *B. burgdorferi* *sensu lato* by adherence of this organism to sphingolipids [86]. Functionally linked to a flagellar protein, HSP-60 is thereby involved in binding to the neural cell surface for intrusion into the CNS [87,88]. As HSP-60 is a major immunodominant antigen of *B. burgdorferi* [89], it comes as no surprise that antibodies to HSP-60 were also detected in the synovial fluid of Lyme arthritis patients [90]. Although still a controversial issue [91–93], molecular mimicry of flagellar epitopes, which are highly antigenic [94], may misdirect antibodies against host tissues as well [95,96]. For pathogens must avoid being destroyed by the immune response while maintaining access to a new host, and protracted antigenic exposure destabilises the immune system.

Molecular, pathologic and microbiological evidence for an involvement of *Borrelia burgdorferi*

Epidemiology cannot replace molecular, experimental and pathological investigation including case reports, despite the fact that these are often dismissed as 'anecdotal'.

Is it possible that the *Borrelia* flagellar basal rod protein (fbrp), implicated in the pathogenesis of sporadic schizophrenia at conception [16,97,98], also plays a pathogenic role in MS following transmission at birth? As fbrp shares an epitope with the human interleukin-1 receptor antagonist IL-1ra (see Table 1 and [99]), it is plausible to assume that such amino acid homology between *B. burgdorferi* and its human host potentially induces and misdirects anti-IL-1ra antibodies. Might re-exposure to the same [100] or similar antigens [101,102] subsequently trigger MS later in life? Conversely, may cross-reacting antibodies acquired against *Treponema* spirochaetes protect migrants

from tropical countries against infection by *Borrelia* spirochaetes and thus antigenic exposure to fbrp?

Not only tropical spastic paraplegia [14] mimics the clinical pattern of MS. In several respects, MS is more reminiscent of neuroborreliosis [102–105], which in its chronic form is supposed to be an autoimmune disease triggered by these spirochaetes [95,105]. MS plaque-derived DNA [106] shows an abundance of transcripts for several heat shock proteins (HSPs), including HSP-70 and anti-HSP-70 antibodies, but apparently not for interleukin-1 (IL-1 β), the underlying pro-inflammatory cytokine (see also [107–110]). This contrasts with other common gram-negative infection, in which HSP-70 induction correlates with elevated levels of IL-1 β transcripts [111]. A putative anti-IL-1ra immune response against IL-1 receptor antagonists (IL-1ra) might therefore explain this specific post-transcriptional dysbalance at the level of the IL-1 receptor unleashing a cascade of ruinous inflammatory cytokines. There are three arguments in support of such a role: reduced genetic expression of IL-1ra versus IL-1 β has been associated with disease severity in MS [112]. IL-1ra can be increased by interferon beta [113], the first neuromodulatory drug approved for the treatment of MS [4]. Conversely, lower IL-1ra versus higher IL-1 activity enhances inflammation, whereas a dysbalance in favour of IL-1ra versus IL-1 reportedly mitigates this reaction in MS, experimental allergic encephalomyelitis (EAE) and *B. burgdorferi* induced Lyme disease [114,115].

When in 1925 Adams et al. [116] inoculated rhesus monkeys with material from MS plaques, spirochaetes emerged in their ventricular fluid after several months. More recently, cystic structures originating from *B. burgdorferi* were found in eight of ten MS patients by immunofluorescence and in all the MS patients by use of transmission electron microscopy and staining after culture [21]. The patients originated from a well-defined coastal area of southern Norway, where Lyme borreliosis

[117,118] as well as MS [12,15] is highly endemic. No such cysts could be observed in the five controls with either method, but the investigators noted a similarity between those found in the MS patients and the cystic forms characteristic of spirochaetes and chronic *B. burgdorferi* infection. More importantly, the cysts of the MS patients exhibited positive reactions to antispirochaetal antiserum [21].

In analogy to the induction of hepatocellular carcinomas upon chronic hepatitis B infection acquired during delivery [24] and in analogy to lymphomas of the skin induced by chronic *B. burgdorferi s.l.* infection [117,118], we would expect an association of MS and ALS with neoplastic transformations of the lymphatic system. A significant correlation between non-Hodgkin's lymphomas and MS appears to exist [119], and in a review of neurolymphomatosis a case was documented with schizophrenia and anterior horn involvement, a hallmark of ALS [120]. As the cause could not be identified, a virally mediated autoimmune pathogenesis was proposed.

Whatever the pathogenic link between schizophrenia and MS - an infection followed by an altered immunologic response [5] or a continuum of chronic inflammatory CNS disorders including neuroborreliosis, syphilis, or viral encephalitis [121] - we would expect an analogous link between schizophrenia and ALS in terms of latitude [53,54], season (see Figure 2) and causality [122].

Among relatives of Ashkenazi immigrants to metropolitan New York suffering from schizophrenia [123], the prevalence of ALS compared to the expected population rate in the USA, where *B. garinii* is non-endemic, was more than a hundred times higher. Not surprisingly after all, the countries of ancestral origin included eastern Europe and Russia [123] areas where *B. garinii* is endemic [117,118].

Up to ten percent of patients initially diagnosed as having ALS are re-diagnosed as having a disease other than ALS [124]. Although it seems unlikely that infection by *B. burgdorferi* is a frequent cause of ALS [117], a discrete subset of patients living in hyperendemic areas appears to be significantly more likely to have immunologic evidence of exposure to *B. burgdorferi* than do controls, and some of these patients do appear to improve if treated with antibiotics [125].

Is the epidemiological association biologically plausible?

Although the nosological criteria for MS and schizophrenia have high diagnostic reliability, affected individuals may differ substantially in the specific profile of signs and symptoms, as well as in the severity and course of their illness. What we recognise clinically as 'schizophrenia' or 'MS', is likely to encompass a complex set of disorders. A

major task of future studies will thus be to resolve the question of heterogeneity in MS [2] as well as its aetiological overlap with other disease processes.

Seroepidemiological studies relating *B. burgdorferi* to MS [126-128] and ALS [125,129,130] have produced conflicting results. However, when entering the CNS, microorganisms can undergo antigenic [131] and extensive metabolic changes, which prevent them from being recognised by the current serologic test methods. These changes protect *B. burgdorferi* from the host's immune system and reduce the effect of antibiotics [21,132]. Although infections all induce specific antibodies and cell-mediated immunity, microbial virulence factors have not often been individually defined or even identified [133]. There are hundreds or thousands of antigens, and immune responses develop to many of these. Resistance to infection, however, depends on the reaction against a few antigens on the surface of the microorganism [133]. The flagellar basal rod protein of *B. burgdorferi* (fbrp) responsible for locomotion, adherence and host cell penetration is part of such a virulence factor. Influencing parasite-host interaction, gram-negative bacteria use this type of basal ring assembly to secrete and translocate flagellar and virulence proteins directly into the cytoplasm of their host cell [134]. While temperature is a key environmental cue for the switch between motility and plasmid-encoded gene expression of virulence, stress-related degradation of the secreted substrates is accordingly prevented by the chaperonising function of HSPs [135]. The coincidence of both genetic and antigenic exposure to *Borrelia* fbrp is therefore not casual, but a highly specific pathogenic event. Mutations by homologous recombination affecting the implanting blastocyst at conception [16] and chronic infection afflicting the immunologically immature newborn upon delivery [24], it is conjectured, will subsequently trigger sporadic congenital schizophrenia and MS respectively.

Conclusion

That maternal infection by *B. burgdorferi s. l.* poses a risk to the neonate cannot be excluded. The global epidemiological clustering by season and locality rather emphasises a causal relation between MS and sporadic schizophrenia, which derives from both genetic and antigenic exposure to a spirochaetal virulence factor at conception and birth. The identification of flagellar *Borrelia* DNA on human CB1 and its relation to IL-1 receptor dysfunction reminds us of Virchow's postulate in 1849: "In searching for pathological systems one must clearly not construct nosological but only etiological ones" [136]. It is hoped that this correlation might encourage a new direction of neuropsychiatric research in molecular epidemiology.

List of abbreviations used

AIDS: acquired immunodeficiency syndrome

ALS: amyotrophic lateral sclerosis

B. burgdorferi s.l.: *Borrelia burgdorferi sensu lato*

CB1: central cannabinoid receptor gene

CNS: central nervous system

DNA: desoxyribonucleic acid

EAE: experimental autoimmune encephalomyelitis

flbp: flagellar basal rod protein

HLA: major histocompatibility complex

HSP: heat shock protein

IL-1: interleukin-1

IL-1ra: interleukin-1 receptor antagonist

MS: multiple sclerosis

PR: prevalence rate

Acknowledgement

To RP re-diagnosed as having a birth defect, MS, ALS, and ultimately atypical neuroborreliosis many thanks for her trust and patience.

References

- Akil M and Weinberger D **Neuropathology and the Neurodevelopmental Model**. In: *The Neuropathology of Schizophrenia* (Edited by: Harrison PJ, Roberts GW) Oxford, University Press 2000, 189-212
- Compston A and Coles A **Multiple sclerosis**. *Lancet* 2002, **359**:1221-1231
- McDonald WI, Ron MA and Giovannoni G **Multiple sclerosis and its pathophysiology**. In: *Diseases of the Nervous System* (Edited by: Asbury AK, McKhann GM, McDonald WI, Goadsby PJ, McArthur JC) Cambridge, University Press 2002, 1606-1619
- Miller DH and Thompson AJ **The diagnosis and management of multiple sclerosis**. In: *Diseases of the Nervous System* (Edited by: Asbury AK, McKhann GM, McDonald WI, Goadsby PJ, McArthur JC) Cambridge, University Press 2002, 1620-1632
- Stevens JR **Schizophrenia and multiple sclerosis**. *Schizophr Bull* 1988, **14**:231-241
- Beaglehole R, Bonita R and Kiehlström T **Basic Epidemiology**. Geneva, World Health Organization Press 1993,
- Templer DI, Cappelletty GG and Kauffman I **Schizophrenia and multiple sclerosis. Distribution in Italy**. *Br J Psychiatry* 1988, **153**:389-390
- Templer DI, Trent NH, Spencer DA, Trent A, Corgiat MD, Mortensen PB and Gorton M **Season of birth in multiple sclerosis**. *Acta Neurol Scand* 1992, **85**:107-109
- Templer DI, Regier MW and Corgiat MD **Similar distribution of schizophrenia and multiple sclerosis**. *J Clin Psychiatry* 1985, **46**:73
- McDonald C and Murray RM **Early and late environmental risk factors for schizophrenia**. *Brain Res Brain Res Rev* 2000, **31**:130-137
- Poser CM **The epidemiology of multiple sclerosis: a general overview**. *Ann Neurol* 1994, **36**(Suppl 2):S180-189
- Kurtzke JF **Epidemiologic evidence for multiple sclerosis as an infection**. *Clin Microbiol Rev* 1993, **6**:382-427
- Marie P **Sclerose en plaque et maladies infectueuses**. *Prog Med* 1884, **12**:287-289
- Hawkes CH **Is multiple sclerosis a sexually transmitted infection?** *J Neurol Neurosurg Psychiatry* 2002, **73**:439-443
- Rosati G **The prevalence of multiple sclerosis in the world: an update**. *Neurol Sci* 2001, **22**:117-139
- Fritzsche M **Seasonal correlation of sporadic schizophrenia to Lyme borreliosis**. *Int J Health Geogr* 2002, **1**:2
- Burkot TR, Schriefer ME and Larsen SA **Cross-reactivity to Borrelia burgdorferi proteins in serum samples from residents of a tropical country nonendemic for Lyme disease**. *J Infect Dis* 1997, **175**:466-469
- Reik L Jr **Neurosyphilis**. In: *Diseases of the Nervous System* (Edited by: Asbury AK, McKhann GM, McDonald WI, Goadsby PJ, McArthur JC) Cambridge, University Press 2002, 1766-1776
- Steiner G **Acute plaques in multiple sclerosis, their pathogenic significance and the role of spirochaetes as an etiological factor**. *J Neuropathol* 1952, **11**:343-372
- Steiner G **Morphology of Spirochaeta myelophtora in multiple sclerosis**. *J Neuropathol* 1954, **13**:221-229
- Brorson O, Brorson SH, Henriksen TH, Skogen PR and Schoyen R **Association between multiple sclerosis and cystic structures in cerebrospinal fluid**. *Infection* 2001, **29**:315-319
- Schlesinger PA, Duray PH, Burke BA, Steere AC and Stillman MT **Maternal-fetal transmission of the Lyme disease spirochete, Borrelia burgdorferi**. *Ann Intern Med* 1985, **103**:67-68
- Silver RM, Yang L, Daynes RA, Branch DW, Salafia CM and Weis JJ **Fetal outcome in murine Lyme disease**. *Infect Immun* 1995, **63**:66-72
- Zinkernagel RM **Maternal antibodies, childhood infections, and autoimmune diseases**. *N Engl J Med* 2001, **345**:1331-1335
- Roach LL, Rosenberg S and Ichelson RR **Immunological consideration of an antigenic fraction from cultures of spirochetes isolated from cerebrospinal fluid of multiple sclerosis cases: preliminary report**. *Am J Med Sci* 1959, **237**:8-11
- Fumarola D **Multiple sclerosis and Borrelia burgdorferi**. *Lancet* 1986, **2**:575
- Stürchler D **Endemic Areas of Tropical Infections**. Toronto, Hans Huber Publishers 1988,
- Torrey EF, Miller J, Rawlings R and Yolken RH **Seasonality of births in schizophrenia and bipolar disorder: a review of the literature**. *Schizophr Res* 1997, **28**:1-38
- Olsen B, Duffy DC, Jaenson TG, Gylfe A, Bonnedahl J and Bergstrom S **Transhemispheric exchange of Lyme disease spirochetes by seabirds**. *J Clin Microbiol* 1995, **33**:3270-3274
- Torrey EF, Miller J, Rawlings R and Yolken RH **Seasonal birth patterns of neurological disorders**. *Neuroepidemiology* 2000, **19**:177-185
- Lauer K **Ecologic studies of multiple sclerosis**. *Neurology* 1997, **49**(2 Suppl 2):S18-26
- Groebke-Lorenz W, Balint S, Frey UP, Giger A, Haner C, Savary B, Studer A, von Werra P and Wüthrich R **Multiple sclerosis in northwestern Switzerland**. *Schweiz Med Wochenschr* 1992, **122**:582-587
- Lauer K and Firnhaber W **Multiple sclerosis mortality 1973-1987 in Baden-Württemberg in comparison with socio-geographical variables**. *Nervenarzt* 1992, **63**:209-212
- Younsi H, Postic D, Baranton G and Bouattour A **High prevalence of Borrelia lusitaniae in Ixodes ricinus ticks in Tunisia**. *Eur J Epidemiol* 2001, **17**:53-56
- Piesman J, Clark KL, Dolan MC, Happ CM and Burkot TR **Geographic survey of vector ticks (Ixodes scapularis and Ixodes pacificus) for infection with the Lyme disease spirochete, Borrelia burgdorferi**. *J Vector Ecol* 1999, **24**:91-98
- Hung TP, Landsborough D and His MS **Multiple sclerosis amongst Chinese in Taiwan**. *J Neurol Sci* 1976, **27**:459-484
- Sonenshine DE **Biology of Ticks**. New York, Oxford University Press 1989,
- McGrath JJ and Welham JL **Season of birth and schizophrenia: a systematic review and meta-analysis of data from the Southern Hemisphere**. *Schizophr Res* 1999, **35**:237-242
- Hudson BJ, Stewart M, Lennox VA, Fukunaga M, Yabuki M, Macorison H and Kitchener-Smith J **Culture-positive Lyme borreliosis**. *Med J Aust* 1998, **168**:500-502

40. Goldstein G, Hall W and Andrews G **The incidence of schizophrenia in New South Wales, Australia. A psychiatric register study.** *Acta Psychiatr Scand* 1984, **70**:220-227
41. Cook SD and Dowling PC **Distemper and multiple sclerosis in Sitka, Alaska.** *Ann Neurol* 1982, **11**:192-194
42. King HO and Bleaney AA **The Falkland Islands morbidity survey.** *J R Coll Gen Pract* 1982, **32**:535-546
43. Fritzsche M and Schmidli J **Seasonal fluctuation in schizophrenia.** *Am J Psychiatry* 2002, **159**:499-500
44. Jensen PM and Frandsen F **Temporal risk assessment for Lyme borreliosis in Denmark.** *Scand J Infect Dis* 2000, **32**:539-544
45. Suvisaari JM, Haukka JK and Lonnqvist JK **Season of birth among patients with schizophrenia and their siblings: evidence for the procreational habits hypothesis.** *Am J Psychiatry* 2001, **158**:754-757
46. Korenberg EI **Seasonal population dynamics of ixodes ticks and tick-borne encephalitis virus.** *Exp Appl Acarol* 2000, **24**:665-681
47. Sadvnick AD and Yee IM **Season of birth in multiple sclerosis.** *Acta Neurol Scand* 1994, **89**:190-191
48. Torrey EF, Torrey BB and Peterson MR **Seasonality of schizophrenic births in the United States.** *Arch Gen Psychiatry* 1977, **34**:1065-1070
49. Clover JR and Lane RS **Evidence implicating nymphal Ixodes pacificus (Acari: ixodidae) in the epidemiology of Lyme disease in California.** *Am J Trop Med Hyg* 1995, **53**:237-240
50. Shaw PJ **Motor neuron disease.** In: *Diseases of the Nervous System* (Edited by: Asbury AK, McKhann GM, McDonald WI, Goadsby PJ, McArthur JC) Cambridge, University Press 2002, 1863-1879
51. Ajdacic-Gross V, Wang J and Gutzwiller F **Season of birth in amyotrophic lateral sclerosis.** *Eur J Epidemiol* 1998, **14**:359-361
52. Charcot J **Lectures on the diseases of the nervous system.** *The New Sydenham Society* 1877, **1**:157-222
53. Annegers JF, Appel S, Lee JR and Perkins P **Incidence and prevalence of amyotrophic lateral sclerosis in Harris County, Texas, 1985-1988.** *Arch Neurol* 1991, **48**:589-593
54. Betemps EJ and Buncher CR **Birthplace as a risk factor in motor neuron disease and Parkinson's disease.** *Int J Epidemiol* 1993, **22**:898-904
55. Poser CM **The dissemination of multiple sclerosis: a Viking saga? A historical essay.** *Ann Neurol* 1994, **36**(Suppl 2):S231-243
56. Brown JS Jr **Geographic correlation of multiple sclerosis with tick-borne diseases.** *Mult Scler* 1996, **1**:257-261
57. Alerstam T **Bird Migration** (Edited by: Christie DA) Cambridge, University Press 1990,
58. Mead C, Ogilvie M, Jackson B, Jackson J, Fullagar P and Oatley T **Birds on the Move** (Edited by: Yelland A, Searchwell M, Ehrlich AM, Ferguson L) London, Harper Collins 1995,
59. Gylfe A, Olsen B, Strasevicius D, Marti Ras N, Weihe P, Noppa L, Ostberg Y, Baranton G and Bergstrom S **Isolation of Lyme disease Borrelia from puffins (Fratercula arctica) and seabird ticks (Ixodes uriae) on the Faeroe Islands.** *J Clin Microbiol* 1999, **37**:890-896
60. Hawkes CH **Twin studies in medicine - what do they tell us?** *QJM* 1997, **90**:311-321
61. Hurko O **Genetics of common neurological disorders.** In: *Diseases of the Nervous System* (Edited by: Asbury AK, McKhann GM, McDonald WI, Goadsby PJ, McArthur JC) Cambridge, University Press 2002, 14-31
62. Helmick CG, Wrigley JM, Zack MM, Bigler WJ, Lehman JL, Janssen RS, Hartwig EC and Witte JJ **Multiple sclerosis in Key West, Florida.** *Am J Epidemiol* 1989, **130**:935-949
63. Anderson JF **Epizootiology of Lyme borreliosis.** *Scand J Infect Dis Suppl* 1991, **77**:23-34
64. Felz MW, Chandler FW Jr, Oliver JH Jr, Rahn DW and Schriefer ME **Solitary erythema migrans in Georgia and South Carolina.** *Arch Dermatol* 1999, **135**:1317-1326
65. Brodmann PA, Suter W, Müller W, Wiedemeier P, Broz P and Bühlmann PJ **Bestandesentwicklung, Aktionsraum und Habitatnutzung der am unteren Zürichsee überwinternden Lachmöwen Larus ridibundus.** *Der Ornithologische Beobachter* 1991, **88**:9-25
66. Gylfe A, Bergström S, Lundström J and Olsen B **Epidemiology; reactivation of Borrelia infection in birds.** *Nature* 2000, **403**:724-725
67. Olsen B, Jaeson T and Bergstrom S **Prevalence of Borrelia burgdorferi sensu lato-infected ticks on migrating birds.** *Appl Env Microbiol* 1995, **61**:3082-3087
68. Salemi G, Ragonese P, Aridon P, Reggio A, Nicoletti A, Buffa D, Conte S and Savettieri G **Is season of birth associated with multiple sclerosis?** *Acta Neurol Scand* 2000, **101**:381-383
69. Stewart GJ **Infection and multiple sclerosis - a new hypothesis?** *J Neurol Neurosurg Psychiatry* 2002, **73**:358-359
70. Vogel F and Motulsky AG **Human Genetics. Problems and Approaches.** Berlin, Springer 1986,
71. Menoret A, Otry C, Labarriere N, Breimer ME, Piller F, Meflah K and Le Pendu J **The expression of carbohydrate blood group antigens correlates with heat resistance.** *J Cell Sci* 1995, **108**:1691-1701
72. Stewart GR, Snewin VA, Walzl G, Hussell T, Tormay P, O'Gaora P, Goyal M, Betts J, Brown IN and Young DB **Overexpression of heat-shock proteins reduces survival of Mycobacterium tuberculosis in the chronic phase of infection.** *Nat Med* 2001, **7**:732-737
73. Hasday JD and Singh IS **Fever and the heat shock response: distinct, partially overlapping processes.** *Cell Stress Chaperones* 2000, **5**:471-480
74. Asea A, Kraeft SK, Kurt-Jones EA, Stevenson MA, Chen LB, Finberg RW, Koo GC and Calderwood SK **HSP70 stimulates cytokine production through a CD14-dependant pathway, demonstrating its dual role as a chaperone and cytokine.** *Nat Med* 2000, **6**:435-442
75. Moseley PL **Heat shock proteins and the inflammatory response.** *Ann N Y Acad Sci* 1998, **856**:206-213
76. Srivastava PK, Menoret A, Basu S, Binder RJ and McQuade KL **Heat shock proteins come of age: primitive functions acquire new roles in an adaptive world.** *Immunity* 1998, **8**:657-665
77. Battistini L, Salvetti M, Ristori G, Falcone M, Raine CS and Brosnan CF **Gamma delta T cell receptor analysis supports a role for HSP 70 selection of lymphocytes in multiple sclerosis lesions.** *Mol Med* 1995, **1**:554-562
78. Leung PS and Gershwin ME **The immunobiology of heat shock proteins.** *J Investig Allergol Clin Immunol* 1991, **1**:23-30
79. Fraser CM, Norris SJ, Weinstock GM, White O, Sutton GG, Dodson R, Gwinn M, Hickey EK, Clayton R, Ketchum KA, Sodergren E, Hardham JM, McLeod MP, Salzberg S, Peterson J, Khalak H, Richardson D, Howell JK, Chidambaram M, Utterback T, McDonald L, Artiach P, Bowman C, Cotton MD and Venter JC **Complete genome sequence of Treponema pallidum, the syphilis spirochete.** *Science* 1998, **281**:375-388
80. Duffell E **Curative power of fever.** *Lancet* 2001, **358**:929-932
81. Hubalek Z, Halouzka J and Heroldova M **Growth temperature ranges of Borrelia burgdorferi sensu lato strains.** *J Med Microbiol* 1998, **47**:929-932
82. Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb JF, Fleischmann RD, Richardson D, Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, van Vugt R, Palmer N, Adams MD, Gocayne J and Venter JC **Genomic sequence of a Lyme disease spirochaete, Borrelia burgdorferi.** *Nature* 1997, **390**:580-586
83. Stamm LV, Gherardini FC, Parrish EA and Moomaw CR **Heat shock response of spirochetes.** *Infect Immun* 1991, **59**:1572-1575
84. Shih CM, Telford SR 3rd and Spielman A **Effect of ambient temperature on competence of deer ticks as hosts for Lyme disease spirochetes.** *J Clin Microbiol* 1995, **33**:958-961
85. Assouf MV, Postic D, Paul G, Nevot P and Baranton G **Western blot analysis of sera from Lyme borreliosis patients according to the genomic species of the Borrelia strains used as antigens.** *Eur J Clin Microbiol Infect Dis* 1993, **12**:261-268
86. Garcia Monco JC, Fernandez Villar B, Rogers RC, Szczepanski A, Wheeler CM and Benach JL **Borrelia burgdorferi and other related spirochetes bind to galactocerebroside.** *Neurology* 1992, **42**:1341-1348
87. Kaneda K, Masuzawa T, Yasugami K, Suzuki T, Suzuki Y and Yanagihara Y **Glycosphingolipid-binding protein of Borrelia burgdorferi sensu lato.** *Infect Immun* 1997, **65**:3180-3185
88. Irani DN and Griffin DE **Host responses in central nervous system infection.** In: *Diseases of the Nervous System* (Edited by: Asbury AK, McKhann GM, McDonald WI, Goadsby PJ, McArthur JC) Cambridge, University Press 2002, 1651-1659

89. Luft GJ, Gorevic PD, Jiang W, Munoz P and Dattwyler RJ **Immunologic and structural characterization of the dominant 66- to 73-kDa antigens of *Borrelia burgdorferi***. *J Immunol* 1991, **146**:2776-2782
90. Mensi N, Webb DR, Turck CW and Peltz GA **Characterization of *Borrelia burgdorferi* proteins reactive with antibodies in synovial fluid of a patient with Lyme arthritis**. *Infect Immun* 1990, **58**:2404-2407
91. Shanafelt MC, Hindersson P, Soderberg C, Mensi N, Turck CW, Webb D, Yssel H and Peltz G **T cell and antibody reactivity with the *Borrelia burgdorferi* 60-kDa heat shock protein in Lyme arthritis**. *J Immunol* 1991, **146**:3985-3992
92. Benoist C and Mathis D **Autoimmunity provoked by infection: how good is the case for T cell epitope mimicry?** *Nat Immunol* 2001, **2**:797-801
93. Schmutzhard E **Multiple sclerosis and Lyme borreliosis**. *Wien Klin Wochenschr* 2002, **114**:539-543
94. Hayashi F, Smith KD, Ozinsky A, Hawn TR, Yi EC, Goodlett DR, Eng JK, Akira S, Underhill DM and Aderem A **The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5**. *Nature* 2001, **410**:1099-1103
95. Aberer E, Brunner C, Suchanek G, Klade H, Barbour A, Stanek G and Lassmann H **Molecular mimicry and Lyme borreliosis: a shared antigenic determinant between *Borrelia burgdorferi* and human tissue**. *Ann Neurol* 1989, **26**:732-737
96. Weigelt W, Schneider T and Lange R **Sequence homology between spirochaete flagellin and human myelin basic protein**. *Immunol Today* 1992, **13**:279-280
97. Fritzsche M **Are cannabinoid receptor knockout mice animal models for schizophrenia?** *Med Hypotheses* 2001, **56**:638-643
98. Fritzsche M **Lateral gene transfer – the missing link between cannabis psychosis and schizophrenia**. *Am J Med Gen* 2002, **114**:512-515
99. **Online Mendelian Inheritance in Man, OMIM (TM). Center for Medical Genetics, Johns Hopkins University (Baltimore, MD) and National Center for Biotechnology Information, Bethesda, National Library of Medicine 2002**.
100. Birner P, Gatterbauer B, Drobna D and Bernheimer H **Molecular mimicry in infectious encephalitis and neuritis: binding of antibodies against infectious agents on Western blots of human nervous tissue**. *J Infect* 2000, **41**:32-38
101. Oldstone MB **Molecular mimicry and autoimmune disease**. *Cell* 1987, **50**:819-820
102. Martin R, Gran B, Zhao Y, Markovic-Plese S, Bielekova B, Marques A, Sung MH, Hemmer B, Simon R, McFarland HF and Pinilla C **Molecular mimicry and antigen-specific T cell responses in multiple sclerosis and chronic CNS Lyme disease**. *J Autoimmun* 2001, **16**:187-192
103. Reik L Jr, Smith L, Khan A and Nelson W **Demyelinating encephalopathy in Lyme disease**. *Neurology* 1985, **35**:267-269
104. Karussis D, Weiner HL and Abramsky O **Multiple sclerosis vs Lyme disease: a case presentation to a discussant and a review of the literature**. *Mult Scler* 1999, **5**:395-402
105. Baig S, Olsson T, Hojberg B and Link H **Cells secreting antibodies to myelin basic protein in cerebrospinal fluid of patients with Lyme neuroborreliosis**. *Neurology* 1991, **41**:581-587
106. Chabas D, Baranzini SE, Mitchell D, Bernard CC, Rittling SR, Denhardt DT, Sobel RA, Lock C, Karpuj M, Pedotti R, Heller R, Oksenberg JR and Steinman L **The influence of the proinflammatory cytokine, osteopontin, on autoimmune demyelinating disease**. *Science* 2001, **294**:1731-1735
107. Dymant DA and Ebers GC **An array of sunshine in multiple sclerosis**. *N Engl J Med* 2002, **347**:1445-1447
108. Whitney LV, Becker KG, Tresser NJ, Caballero-Ramos CI, Munson PJ, Prabhu VV, Trent JM, McFarland HF and Biddison WE **Analysis of gene expression in multiple sclerosis lesions using cDNA microarrays**. *Ann Neurol* 1999, **46**:425-428
109. Lock C, Hermans G, Pedotti R, Brendolan A, Schadt E, Garren H, Langer-Gould A, Strober S, Cannella B, Allard P, Klonowski J, Austin A, Lad N, Kaminski N, Galli SJ, Oksenberg JR, Raine CS, Heller R and Steinman L **Gene-microarray analysis of multiple sclerosis lesions yields new targets validated in autoimmune encephalomyelitis**. *Nat Med* 2002, **8**:500-508
110. Oleksiak MF, Churchill GA and Crawford DL **Variation in gene expression within and among natural populations**. *Nat Genet* 2002, **32**:261-266
111. Huang Q, Liu D, Majewski P, Schulte LC, Korn JM, Young RA, Lander ES and Hacohen N **The plasticity of dendritic cell responses to pathogens and their components**. *Science* 2001, **294**:870-875
112. Schrijver HM, Crusius JB, Uitdehaag BM, Garcia Gonzalez MA, Kostense PJ, Polman CH and Pena AS **Association of interleukin-1 beta and interleukin-1 receptor antagonist genes with disease severity in MS**. *Neurology* 1999, **52**:595-599
113. Sciacca FL, Canal N and Edoardo Grimaldi LM **Induction of IL-1 receptor antagonist by interferon beta: implication for the treatment of multiple sclerosis**. *J Neurovirol* 2000, **6**(Suppl 2):S33-37
114. Porat R, Poutsika DD, Miller LC, Granowitz EV and Dinarello CA **Interleukin-1 (IL-1) receptor blockade reduces endotoxin and *Borrelia burgdorferi*-stimulated IL-8 synthesis in human mononuclear cells**. *FASEB J* 1992, **6**:2482-2486
115. Miller LC, Lynch EA, Isa S, Logan JW, Dinarello CA and Steere AC **Balance of synovial fluid IL-1 beta and IL-1 receptor antagonist and recovery from Lyme arthritis**. *Lancet* 1993, **341**:146-148
116. Adams DK, Blacklock WS and Cluskie JAW **Spirochaetes in ventricular fluid of monkeys inoculated from cases of disseminated sclerosis**. *J Path Bacteriol* 1925, **28**:117-118
117. Satz N **Klinik der Lyme-Borreliose**. Bern, Hans Huber 2002,
118. Gray JS, Kahl O, Lane RS and Stanek G **Lyme Borreliosis: Biology, Epidemiology and Control**. Wallingford, CAB International 2002,
119. Vineis P, Crosignani P, Vigano C, Fontana A, Masala G, Stagnaro E, Miligi L, Costantini AS, Nanni O, Ramazzotti V, Rodella S, Tumino R and Vindigni C **Lymphomas and multiple sclerosis in a multicenter case-control study**. *Epidemiology* 2001, **12**:134-135
120. Diaz-Arrastia R, Younger DS, Hair L, Inghirami G, Hays AP, Knowles DM, Odel JG, Fetell MR, Lovelace RE and Rowland LP **Neurolymphomatosis: a clinicopathologic syndrome re-emerges**. *Neurology* 1992, **42**:1136-1141
121. Engler F and Vetter P **Affective and schizophrenic syndromes in multiple sclerosis. Review of the literature and case reports**. *Schweiz Arch Neurol Psychiatr* 1991, **142**:367-378
122. Howland RH **Schizophrenia and amyotrophic lateral sclerosis**. *Compr Psychiatry* 1990, **31**:327-336
123. Goodman AB **Elevated risks for amyotrophic lateral sclerosis and blood disorders in Ashkenazi schizophrenic pedigrees suggest new candidate genes in schizophrenia**. *Am J Med Genet* 1994, **54**:271-278
124. Traynor BJ, Codd MB, Corr B, Forde C, Frost E and Hardiman O **Amyotrophic lateral sclerosis mimic syndromes: a population-based study**. *Arch Neurol* 2000, **57**:109-113
125. Halperin JJ, Kaplan GP, Brazinsky S, Tsai TF, Cheng T, Ironside A, Wu P, Delfiner J, Golightly M, Brown RH, Dattwyler RJ and Luft BJ **Immunologic reactivity against *Borrelia burgdorferi* in patients with motor neuron disease**. *Arch Neurol* 1990, **47**:586-594
126. Schmutzhard E, Pohl P and Stanek G ***Borrelia burgdorferi* antibodies in patients with relapsing/remitting form and chronic progressive form of multiple sclerosis**. *J Neurol Neurosurg Psychiatry* 1988, **51**:1215-1218
127. Coyle PK, Krupp LB and Doscher C **Significance of reactive Lyme serology in multiple sclerosis**. *Ann Neurol* 1993, **34**:745-747
128. Chmielewska-Badora J, Cisak E and Dutkiewicz J **Lyme borreliosis and multiple sclerosis: any connection? A seroepidemic study**. *Ann Agric Environ Med* 2000, **7**:141-143
129. Waisbren BA, Cashman N, Schell RF and Johnson R ***Borrelia burgdorferi* antibodies and amyotrophic lateral sclerosis**. *Lancet* 1987, **8554**:332-333
130. Mandell H, Steere AC, Reinhardt BN, Yoshinari N, Munsat TL, Brod SA and Clapshaw PA **Lack of antibodies to *Borrelia burgdorferi* in patients with amyotrophic lateral sclerosis**. *N Engl J Med* 1989, **320**:255-256
131. Anguita J, Thomas V, Samanta S, Persinski R, Hernanz C, Barthold SW and Fikrig E ***Borrelia burgdorferi*-induced inflammation facilitates spirochete adaptation and variable major protein-like sequence locus recombination**. *J Immunol* 2001, **167**:3383-3390
132. Brorson O and Brorson SH **An in vitro study of the susceptibility of mobile and cystic forms of *Borrelia burgdorferi* to hydroxychloroquine**. *Int Microbiol* 2002, **5**:25-31
133. Mims C, Nash A and Stephen J **Mims' Pathogenesis of Infectious Disease**. San Diego, Academic Press 2001,

134. Galan JE and Collmer A **Type III secretion machines: bacterial devices for protein delivery into host cells.** *Science* 1999, **284**:1322-1328
135. Page AL and Parsot C **Chaperones of the type III secretion pathway: jacks of all trades.** *Mol Microbiol* 2002, **46**:1-11
136. Virchow R **Die Einheitsbestrebungen in der wissenschaftlichen Medizin.** Berlin, Reimer 1849,

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

